

**Greater New Orleans Surgery Center  
3434 Houma Blvd., Ste 300  
Metairie LA 70006**

---

---

**Operative Report**

Patient Name: \_\_\_\_\_

Date of Procedure: \_\_\_\_\_

Procedure: Epidural Steroid Injection - Levels \_\_\_\_\_  
 Bilateral     Right     Left

Preoperative Diagnosis: \_\_\_\_\_

Post Operative Diagnosis: \_\_\_\_\_

Surgeon: Bernard Landry, M. D.

---

Description of the Procedure:

Pharmacology:	1 cc Sodium Bicarbonate	_____ cc used
	4 cc Lidocaine	_____ cc used
	0.5 cc Isovue M-300	_____ cc used
	3 cc 0.5% Sensorcaine PF	_____ cc used
	1 cc Kenalog (40 mg/ml)	_____ cc used
	3 cc 0.9% NaCl PF	_____ cc used

Procedure: Under fluoroscopic guidance and using a posterolateral approach a 25-gauge needle was advanced into the epidural space thru the translaminar clear space.

Contrast injection confirms the location of the needle tip in the epidural space. The steroid and preservative free anesthetic was then injected without difficulty or complication.

\_\_\_\_\_  
Bernard Landry, M. D.

cc: \_\_\_\_\_